

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76CH03073



PPPL-3572
UC-70

PPPL-3572

Ion Heating by Fast Particle Induced Alfvén Turbulence

D. Gates, N. Gorelenkov, and R.B. White

May 2001



PRINCETON PLASMA PHYSICS LABORATORY
PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY

PPPL Reports Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Availability

This report is posted on the U.S. Department of Energy's Princeton Plasma Physics Laboratory Publications and Reports web site in Calendar Year 2001. The home page for PPPL Reports and Publications is: http://www.pppl.gov/pub_report/

DOE and DOE Contractors can obtain copies of this report from:

U.S. Department of Energy
Office of Scientific and Technical Information
DOE Technical Information Services (DTIS)
P.O. Box 62
Oak Ridge, TN 37831

Telephone: (865) 576-8401
Fax: (865) 576-5728
Email: reports@adonis.osti.gov

This report is available to the general public from:

National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, VA 22161

Telephone: 1-800-553-6847 or
(703) 605-6000
Fax: (703) 321-8547
Internet: <http://www.ntis.gov/ordering.htm>

Ion heating by fast particle induced Alfvén turbulence

D. Gates, N. Gorelenkov, R. B. White

Plasma Physics Laboratory, Princeton University, P.O.Box 451,

Princeton, New Jersey 08543

Abstract

ABSTRACT A novel mechanism that directly transfers energy from Super-Alfvénic energetic ions to thermal ions in high β plasmas is described. The mechanism involves the excitation of compressional Alfvén eigenmodes (CAEs) in the frequency range with $\omega \lesssim \omega_{ci}$. The broadband turbulence resulting from the large number of excited modes causes stochastic diffusion in velocity space, which transfers wave energy to thermal ions. This effect may be important on NSTX [1], and may scale up to reactor scenarios. This has important implications for low aspect ratio reactor concepts, since it potentially allows for the modification of the ignition criterion.

Emissions in the ion cyclotron range of frequency driven by energetic α -particles have been observed in the past on tokamaks [2, 3], but were not observed to have a pronounced effect on the plasma behavior. One of the reasons for this lack of impact was that the energy available to drive the modes is determined by the fraction of energetic particles with a velocity greater than the Alfvén velocity. For the tokamak examples considered, this represents only a small fraction of the total energy available in the alpha particle distribution. If, instead, the fast particle velocity is many times the Alfvén velocity, the power available to sustain the modes can be much larger, on the order of the total fusion power.

The current generation of spherical torus experiments typically operate at very low values of the axial toroidal magnetic field (0.3T to 0.6T). However, the velocity of energetic particles provided by the neutral beam heating sources is quite high. For example on NSTX, which operates with 80keV deuterium neutral beams and 0.3T toroidal field, the neutral beam velocity is typically given by $v_{beam} \sim 4v_A$. The existence of a large class of super-Alfvénic particles can change the regime from one of weakly interacting waves to one where the wave amplitudes become large enough to modify the thermal particle energy. The energy flow is from the fast particles to the waves by

resonance, and from the waves to the thermal particles by stochastic heating.

Ion temperatures greater than the electron temperature have been measured on NSTX during neutral beam heating [4, 5]. Whereas $T_i > T_e$ is not noteworthy in and of itself, the power balance for these discharges is difficult to explain in terms of collisional power balance calculations. $Q_{ie}(\equiv 3/2n\nu_i * (T_i - T_e))$, the power flow from ions to electrons) exceeds the total power going into the ions [6]. The deficit of power is large and, in some cases, on the order of the total neutral beam heating power. The discrepancy is well outside the random errors in the measurements. From this we ascertain that there may be another source of input power to the ions. Another important observation was an off-axis local maximum in the ion temperature. The off-axis peak correlates with the predicted location of the peak amplitude of the compressional Alfvén eigenmode [7]. Also, a series of high frequency plasma modes have been observed during neutral beam heating that have been identified as the compressional Alfvén eigenmode, mentioned above [8]. In addition, a measurement of the lost fast ion flux [9] was 0.1 times the value expected from full particle orbit code predictions. These observations motivate a theoretical consideration of the possible effect of multi-frequency large amplitude CAE modes on ions.

The procedure followed to determine the existence of a wave based ion heating mechanism was to enter the expected wave spectrum from a theoretical model of the CAE mode into a code that calculates particle trajectories. The model used to determine the wave number distribution of the CAE modes is described in [7]. The model calculates which modes are unstable based on a balance between the linear growth rate of the mode and damping on ions and electrons. The primary ion damping mechanism is cyclotron resonance whereas the primary electron damping mechanism is Landau damping.

We note that these modes are largely undamped in the frequency range observed in the experiment $0.2\omega_c < \omega < 0.6\omega_c$. Finite β_e reduces Landau damping for small but finite k_{\parallel} . This can be seen by considering the usual expression for the Landau damping rate, $\gamma_e/\omega = \beta_e\zeta e^{-\zeta^2}$, where $\zeta = \omega/k_{\parallel}v_{the}$. We take the limit $\zeta \ll 1$ and approximate $\omega \sim k_{\perp}v_A$ yielding $\sqrt{m_e/\beta_e m_i} \ll k_{\parallel}/k_{\perp} < 1$. Assuming deuterium ions and $\beta_e = 0.1$ yields $1/19 \ll k_{\parallel}/k_{\perp} < 1$. Finite k_{\parallel} Doppler shifts the modes away from the cyclotron frequency eliminating the resonant thermal ion damping. The lack of damping allows for the modes to grow to sufficient amplitude for the onset of stochasticity.

We assume a cold plasma dielectric tensor for these modes given by:

$$\overline{\overline{\epsilon}}_j = \begin{vmatrix} \epsilon_j & ig_j & 0 \\ -ig_j & \epsilon_j & 0 \\ 0 & 0 & \eta_j \end{vmatrix} \quad \text{with}$$

$$\epsilon_j = \frac{c^2}{v_A^2 \left(1 - \frac{\omega_j^2}{\omega_c^2}\right)}, \quad g_j = \epsilon_j \frac{\omega_j}{\omega_c}, \quad \text{and} \quad \eta_j = -\frac{m_i}{Zm_e} \frac{c^2}{v_A^2} \frac{\omega_c^2}{\omega_j^2}.$$

The dispersion relation is given by:

$$\omega_j^2 = \frac{k_j^2 v_A^2}{2} \left[1 + \cos^2 \theta_j + \sqrt{1 + 2 \left(\frac{2\omega_j^2}{\omega_c^2} - 1 \right) \cos^2 \theta_j + \cos^4 \theta_j} \right]$$

and, for completeness, the wave vectors are:

$$\vec{k}_j = k_{\perp j} (-\sin \phi_j \hat{x} + \cos \phi_j \hat{y}) + k_{\parallel j} \hat{z} = k_{\perp j} \hat{r}_j + k_{\parallel j} \hat{z},$$

$$\vec{E}_{1j} = E_{1j} e^{i\varphi_j} \left[\left(\alpha_j \cos \phi_j - \sin \phi_j \right) \hat{x} + \left(\alpha_j \sin \phi_j + \cos \phi_j \right) \hat{y} - \frac{k_{\parallel j} k_{\perp j} c^2}{\omega_j^2 \eta_j} \hat{z} \right]$$

$$\vec{B}_{1j} = B_{1j} e^{i\varphi_j} \left[\left(\gamma_j \cos \phi_j - \frac{k_{\parallel j}}{k_{\perp j}} \sin \phi_j \right) \hat{x} + \left(\gamma_j \sin \phi_j + \frac{k_{\parallel j}}{k_{\perp j}} \cos \phi_j \right) \hat{y} - \hat{z} \right]$$

$$\text{with} \quad \alpha_j = -i \frac{\omega_c}{\omega_j} \left[1 - \frac{k_{\parallel j}^2 c^2}{\epsilon_j \omega_j^2} \left(1 + \frac{k_{\perp j}^2 c^2}{\omega_j^2 \eta_j} \right) \right], \quad \gamma_j = \frac{k_{\parallel j} k_{\perp j} c^2}{\alpha_j \omega_j^2 \eta_j} - \frac{k_{\parallel j}}{\alpha_j k_{\perp j}},$$

$$\varphi_j = \vec{k}_j \cdot \vec{x} - \omega_j t, \quad B_{1j} = -\frac{\alpha_j k_{\parallel j} c}{\omega_j} E_{1j},$$

$$\theta_j = \cos^{-1} \left(\frac{k_{\parallel j}}{k_j} \right), \quad \text{and} \quad \vec{B}_0 = B_0 \hat{z}.$$

where: the subscript j indexes individual mode frequencies ω_j , subscript 1 indicates the wave field, subscript 0 indicates the background field, B_0

is the equilibrium component of the magnetic field, $v_A = B_0/\sqrt{\mu_0\rho_i}$ is the Alfvén velocity, $\rho_i = m_i n_i$ is the plasma mass density, ω_c is the ion cyclotron frequency, m_e is the electron mass, Z is the charge of the plasma ions, $k_{\perp j}$ and $k_{\parallel j}$ are the perpendicular and parallel wave numbers for the j th mode which define the wavevector \vec{k}_j , ϕ_j is the propagation angle in the x-y plane, θ_j is the angle between \vec{k}_j and the z axis, and r and R are the local minor and major radii of the center of curvature of the flux surface, respectively.

The wave fields calculated in this way are used as input to a code that solves the equation of motion for particles in the prescribed wave combined with a constant background magnetic field. This formulation is the simplest relevant model that can describe the behavior of particles in the presence of a CAE mode spectrum. It has been shown [10] that stochastic heating of ions is possible in the presence of a single large amplitude Alfvén wave below the cyclotron frequency given both a large wave amplitude and oblique propagation, with $k_{\perp} v_A \tilde{B}/(B_0 \omega_c) \gtrsim 0.1$. In the presence of several modes the amplitude threshold can be expected to be much lower.

The actual mode amplitude will be determined by the relative energy transfer efficiencies between the fast ions and the wave, versus between the wave and the thermal ions. Such a theory would require a self-consistent non-

linear treatment of the fast particle distribution function, the wave amplitude and the thermal particle distribution function, which is beyond the scope of the current work. As a first step towards understanding the effects of these modes we simply observe the onset and rate of heating, as a function of a constant mode amplitude.

Because the modes in question will tend to transfer perpendicular energy to the plasma ions, a pitch angle scattering collision operator was included in the equation of motion. This helps avoid the non-physical result of strong anisotropy. The scattering rate was taken to be constant in time with $\nu_i = 0.01\omega_c$. This rate corresponds to an approximate temperature of $\sim 300\text{eV}$ for typical NSTX plasma densities.

The mode spectrum was chosen to roughly match the experimental data. We used $k_\perp = m/r$ with $2 \leq m \leq 8$, and $k_\parallel = n/R$ with $1 \leq n \leq 3$ giving 21 modes. The effective radii, r and R , were chosen so that $0.2\omega_c < \omega < 0.6\omega_c$ and $0.2k_\perp < k_\parallel < 0.5k_\perp$, as indicated by mode stability theory [7]. The propagation angle in the x-y plane, ϕ , was chosen randomly for each mode. The values of k_\parallel were chosen to be in the range that resonates with beam particles, since k_\parallel is unknown.

The results of the calculation are shown in Figures 1 and 2. There is

a clear heating effect that has a strong amplitude dependence. The results were insensitive to most details of the mode spectrum and collision frequency, other than the actual frequency range and the ratio of k_{\perp} to k_{\parallel} . More detailed studies of the heating rate and stochastic thresholds as functions of mode spectrum will be reported in future publications. The self consistent energy balance of these modes will likely be quite complicated, perhaps involving bursting “fishbone” like behavior [11].

The mode amplitudes required for the heating rates in Figure 2 to be similar to that observed on NSTX are on the order of $\sim 2-4$ Gauss in NSTX ($dB_y/B \approx 10^{-3}$), which is plausible since the measured amplitude at the NSTX wall is ~ 4 mGauss. This implies approximately 6-7 e-foldings in the distance between the propagation region and the vessel wall (~ 0.5 m).

We stress that these calculations are only a proof-of-principle and it would be premature to claim agreement with the NSTX data. In order to make a proper comparison between observations and theory a more complete theoretical treatment is required, as well as more detailed experimental data. In particular: the calculation should be done in the correct geometry, the actual growth rates and damping should be used to ascertain the actual expected spectrum, and all modes to higher order in the cyclotron resonance should

be included, with all elements configured to match the experiment.

The result of this calculation is important at many levels. As mentioned above, it is important for the current generation of low aspect ratio toroidal confinement devices, since in these devices the velocity of particles from the neutral beams is well in excess of the Alfvén velocity. It is important to note that the conditions for these modes to grow are substantially easier to arrange at low aspect ratio. The constraints that imply low aspect ratio are that $v_f \gg v_A$, and $\rho_f < a$, where a is the radius of the device and v_f and ρ_f are the relevant fast particle velocity and orbit size. Also one must avoid Landau damping for small but finite k_{\parallel} , as above.

Perhaps more importantly, the velocity of fusion alpha particles in proposed ST reactor concepts, such as ARIES-ST [12], is also well in excess of the Alfvén velocity. In particular, the ARIES-ST design has a peak toroidal field on axis of 2.14T which makes the fusion α -particle velocity $v_{\alpha} \sim 5v_A$. Therefore the CAE modes observed on NSTX could be apparent in a larger ST device with a significant fusion α population. The remaining issue is to determine what fraction of the α energy distribution, which is more isotropic, would interact with the CAE mode. Such a calculation is beyond the scope of this paper.

The implications of coupling fast particle energy directly to ions are covered in detail in [13, 14], and are readily summarized. The criterion for fusion heating power exceeding plasma loss power in a deuterium-tritium reaction is given by $n_i^2 T_i^2 / (n(T_e + T_i) / \tau_E) \gtrsim 10^{21} (\text{keV} \cdot \text{s} \cdot \text{m}^{-3})$. The ignition criterion assumes the primary energy transfer mechanism from fusion α 's is via collisional coupling of the α energy to electrons, which then collisionally couple to the fuel ions. This implies $T_e \gtrsim T_i$, simplifying to the usual criterion $n_i T_i \tau_E \gtrsim 2 \times 10^{21} (\text{keV} \cdot \text{s} \cdot \text{m}^{-3})$. If a direct ion heating effect exists, the higher ion temperature leads to an additional factor f yielding $f n_i T_i \tau_E \gtrsim 2 \times 10^{21} (\text{keV} \cdot \text{s} \cdot \text{m}^{-3})$, where $f = 2 / (1 + T_e / T_i)$. The improvement can be larger if τ_E decreases more strongly with increasing T_e than with increasing T_i .

A concept somewhat similar to the one proposed here, popularly referred to as α -channeling, has been proposed in the past [15]. The mechanism we propose has several advantages over the α -channeling mechanism. Firstly, the modes we discuss occur naturally in the presence of super-Alfvénic particles, therefore excitation power may not be required. Secondly, the mode amplitude heating threshold is small, minimizing concerns about rapid fast particle scattering losses due to large wave amplitudes. Thirdly the waves

extract energy from the fast particles and scatter them into passing orbits, improving the fast particle confinement. Finally, the ion heating mechanism is robust, since the mode frequency is naturally near the ion cyclotron resonance.

A more similar, but less well known, calculation [16] considers the tokamak case with $v \gtrsim v_A$, $\omega \sim 3 - 6\omega_c$ and $k_{\parallel} \simeq 0$. This case shows negligible heating. In particular the $v \gtrsim v_A$ constraint limits the energy available to the modes, the high frequency degrades the coupling to the bulk ions, and the $k_{\parallel} \simeq 0$ condition (which is forced by the low β_e in a tokamak, see above) reduces the doppler shift away from the cyclotron frequency, preventing the modes from growing to sufficient amplitude for stochastic heating to take place. The far from optimum parameter regime eliminates the stochastic heating effect. The insight shown in [16] is nonetheless remarkable.

In conclusion, a novel mechanism for transferring energy directly from a fast ion population to bulk thermal ions in a magnetically confined toroidal plasma with low aspect ratio has been identified. The mechanism involves stochastic heating of thermal particles by Alfvén waves driven by the Super-Alfvénic fast particles. These waves have frequencies such that $\omega \sim \omega_c/2$. This effect is a possible candidate for explaining the higher than expected

ion temperatures seen during neutral beam heating experiments on NSTX. In particular, the off-axis peak in the ion temperature which correlates with the predicted mode location, the experimental observation of the modes, and the lower than expected fast ion loss are strong circumstantial evidence for this effect. If confirmed on NSTX, this effect has a high probability of scaling to an ST reactor. Finally we note that, following the existing experimental observations, we have included only modes with frequencies below the cyclotron frequency. The theory of these modes also predicts resonances at Doppler shifted multiples of the cyclotron frequency. We speculate that modes with frequencies $\omega > \omega_c$ will also contribute to the heating effect, lowering the threshold and/or raising the power transfer to the plasma ions.

The authors would like to acknowledge R. Bell, D. Darrow, E. Fredrickson, S. M. Kaye, B. LeBlanc, J. Menard, and S. Sabbagh for describing the results from NSTX, and D. Mueller and M. Gryaznevich for helpful discussions. This work was supported by the U.S. Department of Energy Grant under contract number DE-AC02-76-CH03073.

References

- [1] M. Ono, et al., Nucl. Fus. **40** (2000) 557
- [2] G. A. Cottrell, R. O. Dendy, Phys. Rev. Lett., **60** (1988) 33
- [3] S. Caufmann, R. Majeski, Rev. Sci. Instrum. **66** (1995) 817
- [4] B. LeBlanc, private communication (2001)
- [5] R. E. Bell, private communication (2001)
- [6] S. M. Kaye, private communication
- [7] N. N. Gorelenkov, C. Z. Cheng, Phys. Plasmas **2** (1995) 1961
- [8] E. Fredrickson, et al., submitted to Phys. Rev. Lett. (2001)
- [9] D. Darrow, private communication
- [10] L. Chen, Z. Lin, R. B. White, Submitted to Phys. Rev. Lett., (2001)
- [11] L. Chen, R. B. White, M. N. Rosenbluth, Phys. Rev. Lett. **52** (1984) 1122
- [12] S. C. Jardin, et al., to be published in J. Fusion Technology, (2001)
- [13] J. F. Clarke, Nucl. Fusion **34** (1980) 1541
- [14] N. J. Fisch and M. C. Herrmann, Nucl. Fusion, **34** (1994) 1541
- [15] N. J. Fisch and M. C. Herrmann, Nucl. Fusion, **35** (1995) 1753
- [16] D. J. Sigmar, Proc. Physics of Plasmas in Thermonuclear Regimes, International School of Plasma Physics, Varenna, Italy, (1979), 237




Figure 1: Ion energy in keV vs time, for $\delta B_y/B = 6.0 \times 10^{-4}$.




Figure 2: Heating rate (keV/s) vs. $\delta B_y/B$.

External Distribution

Plasma Research Laboratory, Australian National University, Australia
Professor I.R. Jones, Flinders University, Australia
Professor João Canalle, Instituto de Fisica DEQ/IF - UERJ, Brazil
Mr. Gerson O. Ludwig, Instituto Nacional de Pesquisas, Brazil
Dr. P.H. Sakanaka, Instituto Fisica, Brazil
The Librarian, Culham Laboratory, England
Library, R61, Rutherford Appleton Laboratory, England
Mrs. S.A. Hutchinson, JET Library, England
Professor M.N. Bussac, Ecole Polytechnique, France
Librarian, Max-Planck-Institut für Plasmaphysik, Germany
Jolan Moldvai, Reports Library, MTA KFKI-ATKI, Hungary
Dr. P. Kaw, Institute for Plasma Research, India
Ms. P.J. Pathak, Librarian, Institute for Plasma Research, India
Ms. Clelia De Palo, Associazione EURATOM-ENEA, Italy
Dr. G. Grosso, Instituto di Fisica del Plasma, Italy
Librarian, Naka Fusion Research Establishment, JAERI, Japan
Library, Plasma Physics Laboratory, Kyoto University, Japan
Research Information Center, National Institute for Fusion Science, Japan
Dr. O. Mitarai, Kyushu Tokai University, Japan
Library, Academia Sinica, Institute of Plasma Physics, People's Republic of China
Shih-Tung Tsai, Institute of Physics, Chinese Academy of Sciences, People's Republic of China
Dr. S. Mirnov, TRINITI, Troitsk, Russian Federation, Russia
Dr. V.S. Strelkov, Kurchatov Institute, Russian Federation, Russia
Professor Peter Lukac, Katedra Fyziky Plazmy MFF UK, Mlynska dolina F-2, Komenskeho
Univerzita, SK-842 15 Bratislava, Slovakia
Dr. G.S. Lee, Korea Basic Science Institute, South Korea
Mr. Dennis Bruggink, Fusion Library, University of Wisconsin, USA
Institute for Plasma Research, University of Maryland, USA
Librarian, Fusion Energy Division, Oak Ridge National Laboratory, USA
Librarian, Institute of Fusion Studies, University of Texas, USA
Librarian, Magnetic Fusion Program, Lawrence Livermore National Laboratory, USA
Library, General Atomics, USA
Plasma Physics Group, Fusion Energy Research Program, University of California at San
Diego, USA
Plasma Physics Library, Columbia University, USA
Alkesh Punjabi, Center for Fusion Research and Training, Hampton University, USA
Dr. W.M. Stacey, Fusion Research Center, Georgia Institute of Technology, USA
Dr. John Willis, U.S. Department of Energy, Office of Fusion Energy Sciences, USA
Mr. Paul H. Wright, Indianapolis, Indiana, USA

The Princeton Plasma Physics Laboratory is operated
by Princeton University under contract
with the U.S. Department of Energy.

Information Services
Princeton Plasma Physics Laboratory
P.O. Box 451
Princeton, NJ 08543

Phone: 609-243-2750
Fax: 609-243-2751
e-mail: pppl_info@pppl.gov
Internet Address: <http://www.pppl.gov>